

ACCEPTED MANUSCRIPT • OPEN ACCESS

The unprecedented coupled ocean-atmosphere summer heatwave in the New Zealand region 2017/18: drivers, mechanisms and impacts

To cite this article before publication: James Salinger *et al* 2019 *Environ. Res. Lett.* in press <https://doi.org/10.1088/1748-9326/ab012a>

Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is © 2018 The Author(s). Published by IOP Publishing Ltd.

As the Version of Record of this article is going to be / has been published on a gold open access basis under a CC BY 3.0 licence, this Accepted Manuscript is available for reuse under a CC BY 3.0 licence immediately.

Everyone is permitted to use all or part of the original content in this article, provided that they adhere to all the terms of the licence <https://creativecommons.org/licenses/by/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions may be required. All third party content is fully copyright protected and is not published on a gold open access basis under a CC BY licence, unless that is specifically stated in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

THE UNPRECEDENTED COUPLED OCEAN-ATMOSPHERE SUMMER HEATWAVE IN THE NEW ZEALAND REGION 2017/18: DRIVERS, MECHANISMS AND IMPACTS

M James Salinger,¹ James Renwick,² Erik Behrens,³ A Brett Mullan,³ Howard J Diamond,⁴ Pascal Sirguey,⁵ Robert O Smith,⁶ Michael C T Trought,⁷ Lisa V Alexander,⁸ Nicolas J Cullen,⁹ B Blair Fitzharris,⁹ Christopher D Hepburn,⁶ Amber K Parker¹⁰ and Phil J Sutton³

¹ Department of Geography and Environmental Studies, University of Haifa, Mt. Carmel, Haifa, 31905, Israel

² School of Geography, Environment and Earth Sciences, Victoria University of Wellington, Wellington New Zealand

³ National Institute of Water and Atmospheric Research, Wellington, New Zealand

⁴ NOAA/Air Resources Laboratory, Silver Spring, Maryland 20910, USA

⁵ National School of Surveying, University of Otago, Dunedin, New Zealand

⁶ Department of Marine Science, University of Otago, Dunedin, New Zealand

⁷ Plant and Food Research, Blenheim, New Zealand

⁸ Climate Change Research Centre and ARC Centre of Excellence for Climate Extremes, UNSW Sydney, Sydney, Australia

⁹ Department of Geography, University of Otago, Dunedin, New Zealand

¹⁰ Department of Wine, Food and Molecular Biosciences, Lincoln University, Lincoln, New Zealand

During austral summer (DJF) 2017/18, the New Zealand region experienced an unprecedented coupled ocean-atmosphere heatwave, covering an area of 4 million km². Regional average air temperature anomalies over land were +2.2°C, and sea surface temperature anomalies reached +3.7°C in the eastern Tasman Sea. This paper discusses the event, including atmospheric and oceanic drivers, the role of anthropogenic warming, and terrestrial and marine impacts. The heatwave was associated with very low wind speeds, reducing upper ocean mixing and allowing heat fluxes from the atmosphere to the ocean to cause substantial warming of the stratified surface layers of the Tasman Sea. The event persisted for the entire austral summer resulting in a 3.8±0.6 km³ loss of glacier ice in the Southern Alps (the largest annual loss in records back to 1962), very early Sauvignon Blanc wine-grape maturation in Marlborough, and major species disruption in marine ecosystems. The dominant driver was positive Southern Annular Mode (SAM) conditions, with a smaller contribution from La Niña. The long-term trend towards positive SAM conditions, a result of stratospheric ozone depletion and greenhouse gas increase, is thought to have contributed through association with more frequent anticyclonic “blocking” conditions in the New Zealand region and a more poleward average latitude for the Southern Ocean storm track. The unprecedented heatwave provides a good analogue for possible mean conditions in the late 21st century. The best match suggests this extreme summer may be typical of average New Zealand summer climate for 2081-2100, under the RCP4.5 or RCP6.0 scenario.

Keywords: Anthropogenic global warming, marine heatwave, atmospheric heatwave, Southern Alps glacier ice volume, wine grapes, marine ecosystems.

Introduction

An increasing number of extremely warm summer heatwaves have occurred since 2000 (Schär *et al* 2004, Sparnocchia *et al* 2006, Barriopedro *et al* 2011, Karl *et al* 2012) in several regions globally. Atmospheric heat wave (AHW) frequency has likely increased in Europe, Asia and Australia since about 1950 (Perkins and Alexander 2013). The number of marine heatwave (MHW) days globally has increased 54% since the early 20th century, based on the definition of Hobday *et al* (2016), with an increase of 0.3-0.9 days per year in the New Zealand region (Oliver *et al* 2018).

AHWs and MHWs are caused by specific atmospheric and oceanic factors or by a combination of both. For AHWs blocking high pressure systems are a predominant driver (Brunner *et al* 2016), while for MHWs oceanic heat advection can be a driver (Oliver *et al.* 2017). The frequency and intensity of such heat extremes are influenced by anthropogenic global warming (AGW; Otto *et al* 2012, Kharin *et al* 2013, Oliver *et al* 2018) with more frequent and more intense heat extremes expected as average temperatures rise. Impacts of such heat extremes are significant and diverse, ranging from coral bleaching (Hughes *et al* 2003), to migration of marine species (Poloczanska *et al* 2016) to human mortality rates (Barriopedro *et al* 2011). Terrestrial impacts of AHWs are very significant as in Europe in 2003 where drought caused an estimated 30% reduction in gross primary production, many large wildfires, and extreme glacier melt in the European Alps (Kosatsky 2005). Individual AHWs and MHWs have been investigated in terms of their characterization and physical drivers (Schär *et al* 2004, Benthuyzen *et al* 2014), marine impacts, and the role of AGW (Oliver *et al* 2018).

This study examines the 2017/18 New Zealand coupled regional AHW and MHW from observations and ocean models. It diagnoses both the atmospheric and oceanic drivers and investigates atmospheric diagnostics. The event is described from observational data sets combined with ocean modelling. The event is discussed as an analogue for projected conditions later in the 21st century (Mullan *et al* 2016).

Methods

Observations of atmosphere and ocean surface temperature

The New Zealand air temperature (NZT) series (Mullan *et al* 2010) was used to calculate mean air temperature anomalies from the 1981-2010 normal. From 1940/41 to 2017/18 climate extremes for TX90p (percentage of days when the daily maximum temperature is above the 90th percentile) and TN90p (percentage of days when the daily minimum temperature is above the 90th percentile) were calculated using the ClimPACT2 software (Alexander and Herold 2015). Eight stations were analysed for the 1934/35 event. Summer days $\geq 25^{\circ}\text{C}$, were analysed for the Virtual Climate Station Network (VCSN), where climate variables, are interpolated onto a 4km by 5km grid over New Zealand (Tait *et al* 2006). Counts of days of $\geq 25^{\circ}\text{C}$ were made for each New Zealand region using VCSN for DJF from 1972/73 to 2017/18, and for station data for the length of record, invariably 45 years or more.

Monthly sea surface temperature (SST) observations were obtained from ERSST version 5 (Huang *et al* 2017), which provides monthly SSTs on a $2^{\circ} \times 2^{\circ}$ latitude/longitude grid from 1854. Monthly SST was averaged over the region 140°E to 150°W and 60°S to the Equator and SST anomalies within this box were calculated with respect to the 1981-2010 period.

Daily SST estimates obtained from the National Oceanic and Atmospheric Administration Optimum Interpolation Sea Surface Temperature (OI SST) V2 product (Reynolds *et al* 2007), provides daily SSTs on a 0.25° latitude/longitude grid spanning September 1981-May 2018. A daily resolution of SST time series for the eastern Tasman Sea was generated by averaging daily SSTs over the region $160\text{-}175^{\circ}\text{E}$ and $36\text{-}48^{\circ}\text{S}$. Daily 9 am measurements of SST since 1953 were obtained from the Portobello Marine Laboratory (PML) time-series, in Otago Harbour, southeast South Island (Greig *et al* 1988). Half-hourly measurements of water temperature at 2 m and 10 m depth spanning summer 2017/18 were obtained on the open coast in a kelp forest 20km north of the Otago Harbour.

MHW definition and metrics

Hobday *et al* (2016) definitions were applied to identify MHWs based on daily SST measurements from the OISSTv2, PML datasets, and hindcast model output. This is when mean SSTs exceed a the 90th percentile for at least 5 consecutive days, and with well-defined start and end times. Daily climatological mean and 90th percentile values were calculated over a 30-year baseline period (September 1981 – August 2011) for each day of the year using all SST data within an 11-day window. The time series were smoothed with a 31-day moving window. The warming during each MHW was characterized using the following metrics: duration (continuous period that SSTs exceed the 90th percentile value), maximum and mean intensity (the maximum and mean daily SST anomaly during each MHW). SST anomalies were calculated relative to the 1981-2011 daily average.

Atmospheric circulation

Monthly mean sea level pressure (MSLP) and 500-hPa geopotential height fields were obtained from the NCEP/NCAR Reanalysis (Kalnay *et al* 1996), and from the ERA-Interim reanalysis (Dee *et al* 2011). Anomalies were calculated relative to a 1981-2010 normal period. Atmospheric circulation patterns were compared using anomaly correlation and root mean-square difference over the region 135°E-140°W, 65°S-25°S. Several indices were used to characterize the circulation. The Southern Oscillation Index (SOI) used was the Troup (1965) method as the standardized anomaly of the MSLP difference between Tahiti and Darwin. The Marshall (2003) Southern Annular Mode (SAM) index was used. For the New Zealand region, two circulation indices, Z1 (west-east) and M1 (south-north airflow), as defined in Trenberth (1976) were used. The frequency of occurrence of weather regimes over New Zealand (Kidson 1994) were compared with the 1981- 2010 normal.

Tropical Cyclone (TC) data for the 2017/18 season were analysed from the Southwest Pacific Enhanced Archive of Tropical Cyclones (SPEARTC, Diamond *et al* 2012), using the normal period of 1981-2010.

Ocean Sub-Surface Temperature

Depth-longitude profiles of sub-surface temperature anomalies were calculated from 1981-2010 from GODAS data (Saha *et al* 2006) averaged between 40°S and 45°S, over the depth range 25 to 600m and longitude range 140°E to 150°W.

Temperatures () for the eastern Tasman Sea (160-172°E, 35-45°S) were calculated using all available Argo data (47-66 profiles per month, Roemmich *et al* 2009). Temperature anomalies were calculated by differencing the Argo profiles and co-located temperature profiles from the CARS 2009 climatology (Ridgway *et al* 2002, based on all available data in the region prior to 2009). The eastern Tasman region was chosen because it was an area of strong SST change during this event and is oceanographically uniform, excluding strong gradients and variability associated with the Subtropical Front to the south, the Tasman Front to the north and the East Australian Current (EAC) to the west. The Argo minus CARS vertical temperature anomalies were regionally averaged, given the uniformity of the region and monthly averaged to increase the signal to noise.

Ocean Modelling

The global ocean model hindcast analysed here is based on the ocean sea-ice component of the UK Earth System Model (Storkey *et al* 2018) and the New Zealand Earth System Model (NZESM; Williams *et al* 2016) using 1°x1° eORCA1 grid and 75 vertical z-levels. Ocean physics have been simulated with the Nucleus for European Modelling of the Ocean (NEMO, Madec 2008) model using the recent released (3.6. stable) code version. Sea-ice has been modelled, using the Community Ice CodE (CICE, Hunke *et al* 2017). Atmospheric boundary conditions to force this ocean-only configuration are based on the atmospheric reanalysis product JRA-55-DO v1.3 (Tsujino *et al* 2018) and the ocean hindcast covers the period from 1958 until February 2018. The fully coupled version of this model (HadGEM3-GC3.1) has been shown to perform well against key metrics in a pre-industrial model setup (Menary *et al* 2018).

1 Mixed layer depths have been defined as depths where the difference to surface density
2 exceeds 0.01 kg/m^3 . Thermocline depths describe the depth where the vertical temperature
3 gradient maximum is located. The anomalies presented in Fig. 3 are computed by subtracting a
4 monthly mean climatology, constructed over the period 1968-2017. The definition of MHW follows
5 Hobday *et al* (2016). The air sea heat fluxes have been computed using bulk formula with JRA-
6 55-DO v1.3 air temperatures and modelled SSTs.

8 *Snow and ice data*

9 The end of summer snowline (EOSS) time series (Chinn *et al* 2012, Willsman *et al* 2018)
10 provides the basis to estimate Southern Alps mountain glacier mass balance from 1977 to 2018.
11 To extend the EOSS for the Southern Alps (EOSS_{Alps}) and for the Tasman Glacier (EOSS_{Tas}),
12 glacier volume changes were calculated for the period 1962-2018 using the methodology
13 described by Chinn *et al* (2012). Independent EOSS_{Tas} measurements from snow surveys of the
14 Tasman Glacier were used to extend the established EOSS_{Alps} record to 1962 (Anderton 1975,
15 Chinn 1968, 1969, 1994), since the two are strongly correlated ($r^2 = 0.91$). Linear regression
16 analysis was used to estimate EOSS_{Alps} for the period 1962-1976, then merged with the
17 documented EOSS_{Alps} record through to 2017 (Willsman *et al* 2018). For 2018, EOSS_{Tas} was
18 assessed using a time series of SENTINEL-2 satellite images from 14 March 2018 (NZDT), then
19 converted to EOSS_{Alps} using the regression relationship described above. An alternative estimate
20 of Southern Alps glacier mass balance comes from glacier-wide albedo observations at Brewster
21 Glacier using Moderate Resolution Imaging Spectroradiometer (MODIS, Sirguey *et al* 2016)
22 imagery, which has been correlated to a record of glacier mass balance (Cullen *et al* 2017). The
23 Brewster Glacier EOSS is strongly correlated with EOSS_{Alps} ($r^2=0.85$).

24 Estimates of water stored as seasonal snow in the South Island for 2017-18 are provided
25 by a conceptual model (SnowSim) originally developed for the electricity industry (Fitzharris and
26 Garr 1995, Kerr 2005). This model calculates water stored as seasonal snow for key hydro
27 generating river catchments and is tuned to their long-term water balance. Past estimates are in
28 general agreement with historical observations of snow back to 1930 (Chinn 1981, Breeze *et al*
29 1986, Fitzharris and Grimmond 1982).

32 *Managed ecosystems: Viticulture*

33 The phenology of biological systems, including grapes, is largely controlled by seasonal
34 temperatures. Grapevine phenology models (GFV – Grapevine Flowering Véraison) are used to
35 estimate seasonal variations in the timing of flowering, the onset of ripening (véraison) and
36 harvest (defined as a fruit sugar concentration of $20 \text{ }^\circ\text{Brix}$). The dates of Marlborough (northeast
37 South Island) Sauvignon blanc flowering, véraison and $20 \text{ }^\circ\text{Brix}$ dates were estimated using the
38 temperature-based GFV model of Parker *et al* 2011 and Parker (2013). The GFV model sums the
39 average daily temperatures (model base temperature of 0°C) from 29th August until the
40 appearance of 50% flowering (GFV F^* [the critical degree-day sum] value of 1282 for Sauvignon
41 Blanc) or 50% véraison (GFV F^* value of 2528 for Sauvignon Blanc).

43 *Climate scenarios*

44 Observations during the extreme heatwave event are compared with future climate scenarios
45 generated by Mullan *et al* (2016), who applied statistical downscaling (Mullan *et al* 2001) to
46 projections from global climate models (GCMs) used in the IPCC Fifth Assessment (Flato *et al*
47 2013). Changes in monthly mean temperatures were calculated for the end-of-century (2081-
48 2100) compared to the baseline period 1986-2005, for four of the IPCC Representative
49 Concentration Pathways (RCPs) described in van Vuuren *et al* (2011). The number of GCMs
50 varied from 18 for RCP6.0 to 41 for the baseline climate and RCP8.5 simulations.

53 **Results**

55 *Characteristics of the 2017/18 New Zealand Heatwave*

56 The coupled ocean-atmosphere heatwave in the New Zealand region during austral summer
57 2017/18 was the most intense recorded in the New Zealand and Tasman Sea regions in 150
58
59
60

1 years of land-surface air temperature records, and ~40 years of satellite-derived sea surface
2 temperature (SST) records (BoM and NIWA, 2018). It covered an area of around 4 million km²,
3 equivalent to that of the Indian sub-continent.

4 For DJF 2017/18, both land-surface air temperatures were 2°C or more above 1981-2010
5 averages over the entire New Zealand region, from 35 to 50°S, 150 to 180°E (Fig 1a, 1c, 1d).
6 There was a huge loss of snow and ice in the Southern Alps (Fig. 4). On land, wine-grape
7 véraison and maturation was very advanced (Fig. 5). Marine environmental impacts saw the
8 appearance of marine species normally found farther north, with impacts on farmed salmon and
9 kelp (Morton 2018, Lewis 2018). There were slightly more named tropical cyclones (TCs) than
10 normal over the southwest Pacific and an increase in extratropical transition of TCs around or
11 towards New Zealand Fig. 2b, 2c), compared to averages reported by Diamond *et al* (2013) and
12 Diamond and Renwick (2015).

13 Land and ocean time series show that the entire New Zealand region was the warmest
14 observed. NZT anomalies were 2.2°C above average (Fig. 1a and Table 2), the warmest on
15 record (Salinger 1979, 1980, Mullan *et al* 2010). Indices of temperature extremes for New
16 Zealand (Fig. 1b and Table 2) show the highest percentage of summer warm days and warm
17 nights above the 90th percentile (35% and 31% respectively) back to 1941. Counts of summer
18 days ≥25°C were the highest recorded, back to 1973. For the Tasman Sea (26°–48°S, 150°–
19 174°E) SSTs were 1.5°C above average (Fig. 1c), the largest anomaly on record. The biggest
20 departures from average occurred in January, with the entire Tasman Sea/New Zealand region
21 having a DJF anomaly of around 2°C (Fig. 1d).

22 Based on both atmospheric and ocean metrics, the summer 2017/18 heatwave was the
23 most intense on record – dating back to 1909 for NZT and 1981 for the satellite SST record. The
24 heatwave over land developed in October and November 2017 where NZT were 1.0 and 1.1°C
25 above average respectively (Fig. 1a). December 2017 was warmer (2.4°C) and the heatwave
26 peaked in January 2018 with an NZT anomaly of 3.2°C, the warmest month ever recorded.
27 During February the event abated (NZT +0.9°C). SSTs followed a similar temporal pattern (Fig.
28 1e-f). Applying a MHW definition (Hobday *et al* 2016) to daily satellite SSTs from the eastern
29 Tasman Sea, the MHW had a duration of 147 days from 14 November 2017 to 9 April 2018
30 (Fig.1g), a maximum intensity of 3.7°C (Fig.1h), and a mean intensity of 2.1°C (not shown, see
31 Methods for a description of the MHW metrics). Daily SSTs from a long-term monitoring site at
32 the PML indicated that summer 2017/18 was the warmest on record, since 1953 (Fig. 1f, 1j). The
33 DJF average temperature (anomaly) at PML was 17.8°C (2.4°C), 0.6°C warmer than the previous
34 record set in 1974/1975. Near-shore surface waters at PML were in an MHW state over four
35 distinct periods from October 2017 to March 2018 (Fig. 1f). Of these, the 28-day period from 22
36 November to 20 December 2017 was the most intense MHW on record at PML, with a maximum
37 (mean) intensity of 5.7°C (Fig.1j) and a mean intensity of 3.4°C (not shown).

38 The only other time a New Zealand coupled ocean-atmosphere heatwave of similar
39 magnitude occurred was in 1934/35. That summer was described at the time as “remarkably
40 warm” (Kidson 1935) with a New Zealand-average mean temperature of 18.5°C, 2.7°C above the
41 climatological normal of the day (1901-1930), and 1.9°C above the 1981-2010 normal (Mullan *et al*
42 2010). TX90p and TN90p were 30 and 29% respectively. There was an accompanying MHW
43 (not analysed by Kidson 1935) which developed in November 1934, became most intense in
44 January 1935, then slowly abated. From ERSSTv5 data (Huang *et al* 2017), SST anomalies were
45 1°C above average for the entire season surrounding New Zealand, and 2°C off the west coast of
46 the South Island (not shown).

47 *Atmospheric and oceanic circulation anomalies*

48 Atmospheric circulation anomalies for the season (Fig. 2a) show a pattern of blocking (higher
49 than normal pressures) to the east and southeast of New Zealand, with negative pressure
50 anomalies to the northwest of New Zealand. Over austral spring 2017 and summer 2017/2018 El
51 Niño/Southern Oscillation (ENSO) was in a weak La Niña phase with a Southern Oscillation Index
52 (SOI) of +0.9 in spring, and +0.0 in summer. This would on average promote a tendency towards
53 northerly quarter airflow anomalies in spring, and northeasterly airflow anomalies over New
54 Zealand in summer (Gordon 1986). The SAM was positive (spring +0.99, summer +1.73)

1 especially in November (+3.2) and January (+2.7). The resultant airflow anomalies (Fig. 2a)
2 match well with the phases of ENSO and the SAM, with the contraction of the southern westerlies
3 towards Antarctica. The regional Trenberth circulation indices are consistent with a prevalence of
4 northeasterly flow for the season, with negative values of both M1 and Z1 indices. . New Zealand
5 weather types show a predominance of the Block regime (observed 54.7%+17.6%), and lack of
6 the Zonal regime (6.6%, -18.5%) for DJF with frequent low wind situations. (Fig. 3a-c). The
7 Trough regime occurrence was near normal (38.6%, +0.8%).

8 TC behaviour during the 2017/18 season was consistent with positive SOI and SAM
9 conditions (Diamond and Renwick 2015). There was a total of eight named TCs for the full
10 season; the 7 late season (Feb-Apr) named TCs over the southwest Pacific (Fig. 2b) were close
11 to the expected number (6) for a late season La Niña period (Diamond *et al* 2013). The season's
12 conditions may have led to an increased extratropical transition of tropical cyclones around or
13 towards New Zealand; such a TC anomaly pattern (Fig. 2c) is consistent with both the La Niña
14 and positive SAM state (Diamond and Renwick 2015).

15 The GODAS sub-surface ocean temperature pattern for DJF for 40-45°S (Fig. 2d)
16 indicates very shallow anomalies to the west of the South Island, with a narrow band down to
17 about 50 m east of the South Island. Argo float measurements (Fig. 2e) averaged over the
18 eastern Tasman Sea confirmed a strong surface warming starting in December, peaking with a
19 3°C mean anomaly over this eastern Tasman region. The anomaly decreased through February-
20 April. The anomaly was shallow, mainly confined to the upper 20 m when it formed but deepened
21 slightly as it was eroded from the surface.

22 *Analogue seasons*

23 A subset of past analogue three-month periods was chosen from both the NCEP and ERA-
24 Interim reanalysis 500hPa anomaly fields. The analogues were chosen based on the highest
25 anomaly correlation and lowest RMS difference (RMSD) across the New Zealand/Tasman Sea
26 region, compared to DJF 2017/18. A total of 10 ERA-Interim and 11 NCEP-I analogues were
27 found which met threshold correlation and RMSD values as documented in Table 1a. Three-
28 month average SOI and SAM indices were calculated for each analogue period and compared to
29 values observed in DJF 2017/18. Both reanalyses produced very similar results in terms of SOI
30 and SAM statistics (Table 1b). All were associated with prominent anticyclonic anomalies of up to
31 60 geopotential metres (gpm) centred to the southeast of New Zealand (not shown). The positive
32 anomalies covered an area from 25-60°S, 155°E-155°W. Compared with the analogue cases in
33 Table 1a (average SAM 1.19, and SOI 0.35) DJF 2017/18 was the 3rd (4th) highest SAM value for
34 the ERA-Interim (NCEP-I) samples. The 95% significance level of SAM and SOI values for these
35 cases is 1.63 and 0.76 respectively, based on the analogue cases in Table 1a. Hence, the SAM
36 value (1.73) for DJF 2017/18 is statistically significant with a p-value of < 0.05, while the SOI
37 value (0.03) was not significant so La Niña conditions likely played a minor role.

38 *Ocean model hindcasts*

39 Global ocean model hindcast results were used to characterize the MHW in time and space. In
40 November 2017 the centre of the MHW was in the southern part of the Tasman Sea, and east to
41 the Chatham Islands whereas by December the entire Tasman Sea and the region south of
42 Chatham Rise were in MHW conditions with intensities exceeding 1°C. In January the MHW
43 weakened but still covered most of the Tasman Sea (Fig. 3d-f). The mixed layer (Fig. 3g-i) and
44 thermocline depth (Fig. 3j) became significantly shallower (~10m and ~40m), due to the decrease
45 in wind speeds over this period (Fig. 3a-c) as suggested by the MSLP anomalies. Wind speeds
46 were around 0.5 ms⁻¹ lower than normal during the season over the Tasman Sea, with
47 climatological values in the order of 6-7 m s⁻¹, reflecting a drop of around 30%. The consequence
48 was anomalous low vertical mixing and surface warming, while the broader oceanic heat content
49 was not affected. Area averages over the Tasman Sea (Fig. 3j) suggest that NDJ wind speeds
50 were a record low, while surface heat fluxes were in the normal range. The subsequent lack of
51 vertical mixing caused the SSTs to exceed previous records. This mechanism contrasts with the
52 2015/16 reported MHW (Oliver *et al* 2017), where horizontal heat advection by the East
53 Australian Current was identified to be the driver.

1 The heat wave during the austral summer period appears to be the result of coupled
2 ocean-atmospheric processes discussed above, with the atmospheric process most likely driven
3 by the intensity of the positive phase of the SAM.
4

5 *Southern Alps Ice Volume and Seasonal Snow*

6 Glacier volume changes in the Southern Alps for the period 1962-2018 were estimated from
7 variations in EOSS indices. The ice volume loss in the Southern Alps in 2017/18 was $3.8 \pm 0.6 \text{ km}^3$
8 water equivalent (w.e.), the largest annual volume loss of permanent ice in the last 57 years (Fig
9 4a). Between 1962 and 2018 the volume of ice in the Southern Alps decreased from an estimated
10 60.5 to 37.3 km^3 w.e., with the 23.2 km^3 w.e. change equal to a 38% loss of ice. The summer
11 heat wave in 2017/2018 was responsible for a 9% reduction in total ice volume in the Southern
12 Alps compared to the previous year (2016/17). The estimate for glacier loss in 2017/2018 was
13 compared to an approach using MODIS observations of surface albedo at Brewster Glacier,
14 which yielded a Southern Alps ice volume loss of $3.1 \pm 0.7 \text{ km}^3$ w.e. MODIS albedo observations
15 revealed the fast and uninterrupted depletion of the winter snowpack compared to the long-term
16 trend (Fig. 4b), leading to an early exposure of glacier ice in mid-December 2017. Surface albedo
17 decayed further (lowest since 2000) and the snowline rose to a level close to or above the altitude
18 of the highest point on the glacier. The minimum albedo corresponded to an annual surface mass
19 balance value of -2.2 m w.e. on Brewster Glacier, which is the most negative surface mass
20 balance value since 1977.
21

22 During the 2017-18 snow year, the SnowSim model (Fig. 4c) showed that the estimated
23 water stored as seasonal snow leading up to August and from mid-December 2018 was the
24 lowest on record. From mid-January it became 'negative', indicating all the seasonal snow had
25 melted, with extraordinary loss of permanent glacier snow and ice. The satellite record
26 corroborates this exceptional loss in snow cover compared to previous years.
27

28 *Managed ecosystems: Viticulture*

29 Sauvignon Blanc flowering, véraison (soluble solids (SS) of 8°Brix) and maturation SS of 20°Brix
30 dates estimated by grapevine phenology (GFV) modelling for four major New Zealand wine-
31 growing regions (Gisborne and Hawke's Bay – both east of North Island, Marlborough and
32 Central Otago – inland southern South Island) showed an advancement in véraison and
33 maturation dates. The Marlborough 2017-18 growing season, particularly from mid-December,
34 experienced temperatures far above the long-term average (Fig. 5a). The estimated dates of
35 véraison and maturation (Fig. 5a) were 2 February and 4 March, compared to the long-term
36 averages (1947-2017) of 15 February and 21 March (advanced by 13 and 17 days respectively).
37 Although 2018 was the warmest season overall and ripening temperatures amongst the warmest
38 since 1947 (Trought *et al* 2016, Fig. 5b), the highest mean ripening temperature occurred in
39 1998. While the GFV model dates for 20°Brix indicated an early maturation, the compressed
40 flowering and an exceptionally successful fruit set and higher than average potential yield
41 experienced during the 2017/18 season delayed the harvest. Above-average bunch numbers are
42 predicted for 2019 reflecting warmer temperatures at bunch initiation.
43

44 *Marine ecology*

45 In southern New Zealand, extensive canopies of the habitat-forming kelp *Macrocystis pyrifera*
46 were unusually absent. The $18\text{-}19^\circ\text{C}$ temperature tolerance threshold of *M. pyrifera* (Hay, 1990)
47 was breached three times at PML (Fig. 1f) and in a nearby open coast kelp forest, a threshold
48 exceeded only seven times since 1953. Periods of high temperatures have been associated with
49 regime shifts and loss of kelp forests elsewhere (Wernberg *et al* 2016). Significant salmon
50 (*Oncorhynchus tshawytscha*) stock mortality was reported in the Marlborough Sounds, the
51 country's most important salmon aquaculture area (Eder 2018) with importation of Atlantic
52 Salmon (*Salmo salar*) for the first time (Hulburt 2018). Out-of-range reports of tropical and warm-
53 temperate fish were also common over the summer (Roy 2018; Lewis 2018). Commercial fishers
54 trawling for snapper (*Pagrus auratus*) in the north of the South Island since 1981 estimated the
55 species had spawned six weeks earlier than normal (Morton 2018). For salmon farms in southern
56
57
58
59
60

1 New Zealand SST was above 17°C for January: only a slight increase in mortalities resulted (J.
2 Swart pers. comm).
3

4 *21st Century Climate Scenarios Comparison*

5 Late 21st century climate change projections for New Zealand representative concentration
6 pathways (RCPs; van Vuuren *et al* 2011) of 2.6, 4.5, 6.0 and 8.5 were compared with DJF
7 2017/18 (Table 2). For example, for Northland, there were 33.4 days (averaged over all VCSN
8 grid-points within the Northland Regional Council boundary), compared to 20.4 days in the 1981-
9 2010 climatology, a difference of 64% above climatology.

10 For the scenario projections (columns 2 and 9-12), comparisons use the IPCC
11 convention and compare 2081-2100 relative to 1986-2005. The statistical downscaling procedure
12 (Mullan *et al* 2001) calculates a future change for the mean temperature only; this offset has been
13 applied to the VCSN daily maximum temperatures over 1986-2005, and then exceedances of the
14 25°C threshold counted (following Table 8 of Mullan *et al* 2016). Changes are expressed as the
15 percentage increase of the 2081-2100 count above the 1986-2005 count.

16 The DJF2017/18, counts of days $\geq 25^{\circ}\text{C}$ compare best with end-of-century projections
17 under RCP4.5 or RCP6.0. Changes in observed number of days $\geq 25^{\circ}\text{C}$ ranged from a 10%
18 increase relative to climatology in Gisborne (eastern North Island) to a 195% increase in Taranaki
19 (western North Island). In terms of the climate scenario changes, summer extremes are mostly
20 too small under RCP2.6 and far exceed 2017/18 observations by the end of the 21st century
21 under RCP8.5. Summer temperature departures for 2017/18 compared best with end-of-century
22 changes under RCP4.5 or RCP6.0, where mean annual temperatures increase by 1.4°C or 1.8°C,
23 respectively by 2081-2100 (Table 6, Mullan *et al* 2016).

24 We recognize that 2017/18 is a single-season transient extreme event, whereas
25 conditions in the 2090s (2081-2100 average) under future scenarios are closer to a near-
26 equilibrium state. The scenario changes in managed ecosystems and marine ecology are
27 expected to be in the same direction as seen in 2017/18, but the changes could be larger if the
28 drivers are consistent year after year. Larger changes are likely for slow-response components of
29 the New Zealand climate system, such as the South Island glaciers.
30
31
32

33 **Discussion and Conclusions**

34
35 Most global land areas have experienced more heatwaves since the mid-20th century (Donat *et al*
36 2013). Only recently have MHWs been assessed (Oliver *et al* 2018) and very few studies have
37 examined coupled ocean-atmosphere heatwaves, one such example being that of Olita *et al*
38 (2007) of the Mediterranean in 2003. The unprecedented heatwave in the 2017/18 New Zealand
39 summer (DJF) was such a combined AHW/MHW event. It was the most intense on record with a
40 maximum AHW anomaly of +3.2°C, and MHW anomaly of +3.7°C. The heatwave covered all the
41 land area, the entire central and south Tasman Sea and across to 180°E in the southwest Pacific
42 Ocean, an area of 4 million km². The event was discernible in monitored surface air
43 temperatures, daily remotely-sensed SSTs, nearshore temperature loggers, GODAS and Argo
44 float measurements. Slightly more named TCs than normal occurred in the southwest Pacific,
45 with an above average number of extratropical transitions towards and around New Zealand. The
46 ocean modelling analyses suggest, that a general lack of vertical mixing due to exceptional low
47 winds during this period was the main driver for this MHW event, while earlier studies found that
48 heat advection have triggered previous events in the Tasman Sea (Oliver *et al* 2017).
49

50 Likely climate drivers are depicted qualitatively in Fig. 6, with the role of AGW highlighted.
51 Recent research using climate models by Perkins-Kirkpatrick *et al* (2018) suggests that the
52 2017/18 MHW would have been “virtually impossible” without anthropogenic influence. However,
53 the statistics from the 1934/35 event indicate it has occurred in the past in the observed record
54 and was only 0.3°C cooler. If AGW is adjusted for, it was in effect 0.2°C warmer. Projected
55 changes of pressure and wind for the late 21st century from climate models (Mullan *et al* 2016)
56 show MSLP increases in the DJF period, especially to the south-east of New Zealand. The airflow
57 becomes more north-easterly, and at the same time associated with more (possibly blocking)
58
59
60

1 anticyclones. AGW also produces a trend towards the SAM being more positive resulting in
2 higher MSLPs in the New Zealand region, and a contraction of the southern westerlies poleward,
3 however this trend will likely be countered over the coming several decades by stratospheric
4 ozone recovery (Arblaster *et al* 2011).

5 The New Zealand DJF heatwave shows all these features (Fig. 6), with increased MSLP
6 mainly from the SAM together with the oceanic and atmospheric drivers of ENSO in a weak La
7 Niña phase. This produced very prominent positive height anomalies at the 500-hPa level with
8 positive anomalies of up to 60 gpm centred to the southeast of New Zealand. The atmospheric
9 circulation anomalies were associated with very low wind speeds, around 2 ms⁻¹ lower than
10 normal during the season. Ocean model simulations suggest this caused anomalous low vertical
11 mixing and initiated ocean surface warming, although the broader oceanic heat content was not
12 affected. While surface heat fluxes were normal, the lack of vertical mixing caused the SSTs to
13 exceed previous records and led to increased temperature anomalies.

14 The heatwave was over land and oceanic areas, with significant impacts on the terrestrial
15 environment and marine ecosystems. Tasman glacier snowline estimates show the highest
16 EOSS_{Tas} on record. An estimated record 3.8±0.6 km³ w.e. loss of ice occurred, the largest
17 estimated annual volume loss of permanent ice in the Southern Alps for the last 57 years and a
18 9% reduction compared to the previous year. Seasonal snow storage leading up to August and
19 from mid-December 2018 was the lowest on record. From mid-January it became “negative”,
20 indicating all seasonal snow and some long-term ice had melted. For Marlborough Sauvignon
21 Blanc wine grapes the 2018 season proved to be amongst the warmest since 1947, with véraison
22 and maturation dates advanced by 13 and 17 days respectively. Unprecedented breaches of
23 physiological temperature thresholds for habitat forming kelp and observations of out-of-range
24 occurrence of tropical and warm temperate fish species suggest broader impacts on marine
25 ecosystems.

26 Finally, we have examined the intensity of the warming, and extreme surface air
27 temperatures to provide a future analogue for summer climate scenarios for 2081-2100 for impact
28 studies. The statistics of the anomalies and extremes match well with the RCP4.5 or RCP6.0
29 summer climate projections for the New Zealand region.

30 Our approach has been applied in near-real time as the event developed in the oceanic
31 Southern Hemisphere. This contrasts with previous studies, which have separately examined
32 AHW and MHW events (Sparnocchia *et al* 2006, Barriopedro *et al* 2011), and their physical
33 drivers and mechanisms (Oliver *et al* 2018), and impacts (Benthuyssen *et al* 2014). This also
34 provides a useful analogue for future AGW in the region. In doing so we utilized ocean modelling
35 to diagnose physical drivers and pioneered the use of satellite remote sensing to track glacier ice
36 volume loss for the Southern Alps. We note the importance of long-term observation records as
37 well as remote-sensing observing systems to track the evolution of both AHW and MHW events.
38 This multidisciplinary work has shown that regional heatwaves can develop rapidly and have
39 widespread impacts on ecosystems. It would be valuable for the approach taken here to utilize
40 both atmospheric and oceanic seasonal to interannual climate forecasting systems (Salinger *et al*
41 2016) to become part of a scheme that could provide alerts on the emergence and evolution of
42 regional heatwaves. Under the more extreme RCP8.5 scenario, the observed 2017/18 heatwave
43 anomalies would become typical much earlier in the century.

44 Acknowledgements

45 Updates of the Tasman Glacier end of summer snow line data to 1976 were provided by Dr
46 Trevor Chinn. The contribution of Dr Elizabeth (Betty) Batham is recognized in establishing the
47 long-term PML temperature record. Thanks to Dr Doug Mackie for providing the daily PML SST
48 record used here. Observations from a salmon farm in southern New Zealand were provided by
49 Jaco Swart, Salmon Farm Manager, Sanfords. We also thank our many colleagues and
50 volunteers who have maintained and contributed to the New Zealand marine and climate data
51 sets utilised in this study. Thanks to Ms. Noga Yoselevich for preparing the figures. The sub-
52 surface SST data was collected as part of the MBIE funded Coastal Acidification: Rate, Impacts
53
54
55
56
57
58
59
60

1 and Management (CARIM) project, provided by Kim Currie. This project obtained support through
2 the Deep South National Science Challenge.
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Accepted Manuscript

Table 1a. Detailed 500-hPa Analogue Results by Season [correlation threshold minimum 0.65; RMSE threshold maximum 19 gpm]. These are the results of an analysis of the atmospheric circulation patterns were compared using anomaly correlation and root mean-square difference over the region 135°E-140°W, 65°S-25°S compared to the DJF 2017/18 season (SAM value of 1.73; SOI value of 0.03). The DJF 2017/18 SAM values were statistically significant (p , 0.05) for both ERA-Interim and NCEP I analyses. Bolded text below indicates values are $p < 0.05$.

ERA-Interim	SAM Value	SOI Value	NCEP I	SAM Value	SOI Value
JJA 1979	2.35	-0.20	NDJ 1958	0.90	-0.67
OND 1981	1.39	0.03	DJF1962	1.39	1.17
JJA 1985	0.47	-0.07	JAS 1969	1.50	0.73
DJF 1994	1.54	0.00	JJA1979	2.35	-0.20
FMA 1999	0.54	1.17	JJA 1983	0.70	-0.27
MAM 1999	1.19	0.97	JFM 1999	0.48	1.10
JAS 2005	0.13	-0.10	FMA 1999	0.54	1.17
DJF 2008	2.15	1.57	JAS 2005	0.13	-0.10
DJF 2013	0.88	-0.37	DJF 2008	2.15	1.57
JFM 2018	1.24	0.47	FMA 2013	1.69	0.30
-	-	-	JFM 2018	1.24	0.47

Table 1b. Summary table of SAM index and SOI values and related statistics for close analogue 500 hPa height anomalies from ERA-Interim and NCEP I Reanalysis datasets as documented in Table 1a.

Variable	ERA-Interim (1979-2018)	NCEP I Reanalysis (1948-2018)
# of Overlapping Cases	10	11
Mean SAM Value	1.19	1.19
Mean SOI Value	0.35	0.48
SAM at $p < .05$ level	1.55	1.73
SOI at $p < .05$ level	0.79	0.87
Ranking of DJF 2017/18 SAM	3rd Highest	3rd Highest

Table 2. Statistics of summer temperature percentiles and extremes for the 2017/18 summer and for the RCPs for 2081–2100, shown separately for each New Zealand Regional Council region (following Table 6 of Mullan et al, 2016). Column 2 shows the model ensemble-average change in mean temperature and (in brackets) the 5th to 95th percentile range over 18 to 41 models (dependent on RCP). Columns 3 & 4 show the percentage of time the 2017/18 daily maximum and minimum temperatures, respectively, were above the 1981-2010 90th percentiles. Column 5 gives the observed 2017/18 anomaly relative to 1981-2010. Columns 6-8 show days $\geq 25^{\circ}\text{C}$ from the VCSN for 2017/18 (column 6), 1981-2010 climatology (column 7), and percentage difference (column 8). Columns 9-12 show days $\geq 25^{\circ}\text{C}$ for the scenarios, expressed as the percentage change of 2081-2100 days relative to 1986-2005 days, for four RCPs (2.6, 4.5, 6.0 and 8.5) respectively. The map of regional council areas can be found at: <http://www.localcouncils.govt.nz/>

Region	Summer 2081-2100 change 1986-2005 ($^{\circ}\text{C}$)	2017/18 %TX90p	2017/18 %TN90p	2017/18 mean temp. anomaly $^{\circ}\text{C}$	2017/18 Days $\geq 25^{\circ}\text{C}$ VCSN	Average days $\geq 25^{\circ}\text{C}$ VCSN	2017/18 Diff. %	RCP 2.6 summer change %	RCP 4.5 summer change %	RCP 6.0 summer change %	RCP 8.5 summer change %
Northland rcp 8.5 rcp 6.0 rcp 4.5 rcp 2.6	3.3 (2.4, 5.0) 2.0 (1.4, 3.3) 1.5 (0.9, 2.5) 0.7 (0.4, 1.3)	37	27	1.7	33.4	20.4	64%	52%	125%	175%	305%
Auckland rcp 8.5 rcp 6.0 rcp 4.5 rcp 2.6	3.3 (2.3, 5.1) 2.0 (1.2, 3.6) 1.5 (0.8, 2.6) 0.7 (0.2, 1.4)	46	32	2.0	42.1	17.0	148%	56%	142%	201%	355%
Waikato rcp 8.5 rcp 6.0 rcp 4.5 rcp 2.6	3.3 (2.2, 5.3) 2.0 (1.1, 3.8) 1.5 (0.7, 2.7) 0.7 (0.2, 1.4)	44	9	2.0	39.2	20.5	91%	41%	103%	144%	256%
Bay of Plenty rcp 8.5 rcp 6.0 rcp 4.5 rcp 2.6	3.3 (2.3, 5.0) 2.0 (1.2, 3.6) 1.5 (0.8, 2.6) 0.7 (0.2, 1.3)	24	40	2.0	25.6	14.5	77%	54%	137%	197%	364%
Taranaki rcp 8.5 rcp 6.0 rcp 4.5 rcp 2.6	3.3 (2.2, 5.2) 1.9 (1.0, 3.9) 1.5 (0.7, 2.7) 0.6 (0.1, 1.4)	52	35	2.4	17.1	5.8	195%	71%	203%	309%	634%
Manawatu-Whanganui rcp 8.5 rcp 6.0 rcp 4.5 rcp 2.6	3.3 (2.3, 4.7) 1.9 (1.0, 3.6) 1.5 (0.7, 2.7) 0.7 (0.1, 1.4)	46	38	2.4	34.1	15.5	120%	38%	97%	138%	252%
Gisborne rcp 8.5 rcp 6.0 rcp 4.5 rcp 2.6	3.2 (2.3, 4.6) 2.0 (1.2, 3.4) 1.5 (0.9, 2.5) 0.7 (0.3, 1.3)	20	20	1.8	21.1	19.1	10%	33%	80%	113%	212%
Hawke's Bay rcp 8.5 rcp 6.0 rcp 4.5 rcp 2.6	3.1 (2.3, 4.5) 1.9 (1.2, 3.5) 1.4 (0.8, 2.6) 0.7 (0.3, 1.3)	22	29	2.2	35.3	22.3	58%	31%	73%	101%	184%

References

- Alexander L V A and Herold N 2015 ClimPACTv2 Indices and Software. A document prepared on behalf of the Commission for Climatology (CCI) Expert Team on Sector-Specific Climate Indices (ET-SCI) ftp://ftp.cimh.edu.bb/CariCOF/CLIMPACT/climpact2-master/climpact2-master/ClimPACTv2_manual.pdf
- Ainge Roy E 2018. New Zealand 'marine heatwave' brings tropical fish from 3,000km away. *The Guardian*. Retrieved from www.theguardian.com
- Anderton P W 1975 Tasman Glacier 1971-73. *Hydrological Research: Annual Report No. 33*. Ministry of Works and Development for the National Water and Soil Conservation Organisation, Wellington, New Zealand. 28pp
- Arblaster, J. M., Meehl, G. A., & Karoly, D. J. 2011. Future climate change in the Southern Hemisphere: Competing effects of ozone and greenhouse gases. *Geophysical Research Letters*, 38(2). doi:10.1029/2010GL045384
- Barriopedro D E M, Fischer J, Luterbacher R M. Trigo and García-Herrera R 2011 The hot summer of 2010: Redrawing the temperature record map of Europe *Science* **332** (6026) 220-224
- Benthuyzen J, Feng M and Zhong L 2014 Spatial patterns of warming off Western Australia during the 2011 Ningaloo Nino: quantifying impacts of remote and local forcing *Cont. Shelf Res.* **91** 232–246
- Bowen M, Markham J, Sutton P, Zhang X, Wu Q, Shears *et al* 2017 Interannual Variability of Sea Surface Temperature in the Southwest Pacific and the Role of Ocean Dynamics *J. Climate* **30**, 7481-7492 <https://doi.org/10.1175/JCLI-D-16-0852.1>
- BoM and NIWA 2018 Special climate statement – record warmth in the Tasman Sea, New Zealand and Tasmania. Australian Bureau of Meteorology Special Climate Statement 64, Australian Bureau of Meteorology and NIWA Special Climate Statement, 16pp. <http://www.bom.gov.au/climate/current/statements/scs64.pdf>
- Breeze E D and 8 others 1986 A history of snow avalanches in New Zealand. New Zealand Mountain Safety Council. Avalanche Report 5
- Brunner L, Hegerl G Steiner A K 2017 Connecting atmospheric blocking to European temperature extremes in spring, *J. Climate* **30(2)** 585–594, doi:10.1175/JCLI-D-16-0518.1
- Chinn T 1968 Snow surveys on the Tasman Glacier. *New Zealand Alpine Journal* **22(2)** 411-416
- Chinn T J 1969 Snow Survey Techniques, Waitaki Catchment, South Canterbury. *Journal of Hydrology (New Zealand)* **8(2)** 68-76
- Chinn T J 1981 Snowfall variations, hazards and snow melt. In Mountain Lands Workshop. Christchurch. Ministry of Works and Development. Water and Soil Science Centre, 1-21. (Report WS 525.)
- Chinn T J 1994 Snow and ice balance measurements from the Tasman Glacier, Waitaki Catchment, New Zealand. Client Report 413399.22 Institute of Geological and Nuclear Sciences Ltd, Dunedin, NZ
- Chinn T J, Fitzharris B B, Salinger M J, Willsman A 2012 Annual ice volume changes 1976–2008 for the New Zealand Southern Alps *Glob. Planet. Change* doi:10.1016/j.gloplacha.2012.04.002
- Cullen N J, Anderson B, Sirguey P, Stumm D, Mackintosh A, Conway J P, Horgan H J, Dadic R, Fitzsimons S J, and Lorrey A 2017 An eleven-year record of mass balance of Brewster Glacier, New Zealand, determined using a geostatistical approach *J. Glaciology* **63, (238)** 199-217 doi:10.1017/jog.2016.128
- Dee D P, Uppala S M, Simmons A J, Berrisford P, Poli P, Kobayashi S, Andrae U, Balmaseda M A, Balsamo G, Bauer P, Bechtold P, Beljaars A C M, van de Berg L, Bidlot J, Bormann N, Delsol C, Dragani R, Fuentes M, Geer A J, Haimberger L, Healy S B, Hersbach H, Hólm E V, Isaksen L, Kállberg P, Köhler M, Matricardi M, McNally A P, Monge-Sanz B M, Morcrette J-J, Park B-K, Peubey C, de Rosnay P, Tavolato C, Thépaut J-N, Vitart F 2011 The ERA-Interim reanalysis: configuration and performance of the data assimilation system *Q. J. R. Meteorol. Soc.* **137**: 553–597
- Diamond H J, Lorrey A M and Renwick J A 2013 A Southwest Pacific Tropical Cyclone Climatology and Linkages to the El Niño–Southern Oscillation. *J. Clim.* **26** 3-25 doi: <https://doi.org/10.1175/JCLI-D-12-00077.1>
- Diamond H J and Renwick J A 2015 The climatological relationship between tropical cyclones in the southwest Pacific and the southern annular mod, *Int. J. Climatol.* **35** 613–623
- Diamond H J, Lorrey A M, Knapp K R and Levinson D H 2012. Development of an enhanced tropical cyclone tracks database for the southwest Pacific from 1840-2011. *International Journal of Climatology* **32**: 2240–2250. DOI:10.1002/joc.2412
- Donat M G, Alexander L V, Yang H, Durre I, Vose R, Dunn R J H, Willett K M, Aguilar E, Brunet M, Caesar J, Hewitson B, Jack B, Klein Tank A M G, Kruger A C, Marengo J, Peterson T C, Renom M, Oria Rojas C, Rusticucci M, Salinger M J, Sanhoury Elayah A, Sekele S S, Srivastava A K, Trewin B, Villarreal C, Vincent L A, Zhai P, Zhang X, Kitching S 2013 Updated analyses of temperature and

- precipitation extreme indices since the beginning of the twentieth century: The HadEX2 dataset *J. Geophys. Res. Atmospheres* **118** 2098-2188 DOI: 10.1002/jgrd.50150
- Eder J 2018 Hotter-than-normal water kills off salmon in the Sounds. *Marlborough Express*. Retrieved from www.stuff.co.nz
- Fitzharris B and Garr G E 1995 Simulation of past variability in seasonal snow in the Southern Alps, New Zealand. *Annals of Glaciology* **21** 377-382
- Fitzharris B B and Grimmond C S B 1982 Assessing snow storage and melt in a New Zealand mountain environment. *International Association of Hydrological Sciences Publication* (Symposium at Exeter 1982 — Hydrological Aspects of Alpine and High-Mountain Areas) 161-168.
- Flato G, Marotzke J, Abiodun B, Braconnot P, Chou S C, Collins W, Cox P, Driouech F, Emori S, Eyring V, Forest C, Gleckler P, Guilyardi E, Jakob C, Kattsov V, Reason C, Rummukainen M 2013 Evaluation of Climate Models. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker T F, Qin D, Plattner G K, Tignor M, Allen S K, Boschung J, Nauels A, Xia Y, Bex V and Midgley P M (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA
- Gordon N D 1986 The Southern Oscillation and New Zealand weather *Mon. Wea. Rev.*, **114**, 371-387
- Greig M J, Ridgway N M, Shakespeare B S 1988 Sea surface temperature variations at coastal sites around New Zealand *NZ J. Mar. Freshw. Res.* **22** 391-400
- Hay C H 1990 The distribution of *Macrocyctis* (Phaeophyta: Laminariales) as a biological indicator of cool sea surface temperature, with special reference to New Zealand waters *J. Roy. Soc. NZ* **20**:4 313-336 DOI: [10.1080/03036758.1990.10426716](https://doi.org/10.1080/03036758.1990.10426716)
- Hobday A, Alexander L V, Perkins S E, Smale D A, Straub S C, Oliver E C J, Benthuyzen J A, Burrows M T, Donat M G, Feng M, Holbrook N J, Moore P J, Scannell H A, Sen Gupta A, Wernberg T 2016 A hierarchical approach to defining marine heatwaves *Prog. Oceanogr.* **141** 227-238
- Huang B, Thorne P W, Banzona V F, Boyer T, Cheperin G, Lawrimore J W, Menne M J, Smith T M, Vose R S and Zhang H-M 2017 Extended Reconstructed Sea Surface Temperature version 5 (ERSSTv5), Upgrades, validations, and intercomparisons *J. Climate* **30** 8179-8205, [doi: 10.1175/JCLI-D-16-0836.1](https://doi.org/10.1175/JCLI-D-16-0836.1)
- Hulburt P 2018 Salmon industry in short supply turns to Atlantic seas for help. The Marlborough Express. Retrieved from www.stuff.co.nz
- Hughes T P, Baird A H, Bellwood D R, Card M, Connolly S R, Folke C, Grosberg R, Hoegh-Guldberg O, Jackson J B C, Kleypas J, Lough J M, Marshall P, Nyström M, Palumbi S R, Pandolfi J M, Rosen B, Roughgarden J et al 2003 Climate change, human impacts, and the resilience of coral reefs, *Science* (**80-**) **301** (5635), 929-933, [doi:10.1126/science.1085046](https://doi.org/10.1126/science.1085046).
- Hunke E, Lipscomb W, Jones P, Turner A, Jeffery N and Elliott S 2017 CICE, The Los Alamos Sea Ice Model. <https://www.osti.gov/biblio/1364126>
- Karl T R, Gleason B E, Menne M J, McMahon J R, Helm R R Jr, Brewer M J, Kundel K E, Arndt D S, Privette J L, Bates J J, Groisman P Y and Easterling D R 2012 U. S. Temperatures and drought: recent anomalies and trends *Eos, Trans. Amer. Geo. Union* **93** 473-474 [doi: 10.1029/2012EO470001](https://doi.org/10.1029/2012EO470001)
- Kalnay E, Kanamitsu M, Kistler R, Collins W, Deaven D, Gandin L, Iredell M, Saha S, White G, Woollen J, Zhu Y, Chelliah M, Ebisuzaki W, Higgins W, Janowiak J, Mo K C, Ropelewski C, Wang J, Leetmaa A, Reynolds R, Jenne R, Joseph D 1996 The NCEP/NCAR 40-year reanalysis project, *Bull. Amer. Meteor. Soc.* **77** 437-470
- Kerr T 2005 Snow storage modeling in the Lake Pukaki catchment, New Zealand: an investigation of enhancements to the SNOWSIM model Master's thesis, University of Canterbury, Christchurch, New Zealand
- Kharin V V, Zwiers F W, Zhang X and Wehner M 2013 Changes in temperature and precipitation extremes in the CMIP5 ensemble *Climatic Change* **119** 345-357
- Kidson E 1935 The summer of 1934/35 in New Zealand *NZ Met. Office Note* 16 12pp
- Kidson J W 1994 An automated procedure for the identification of synoptic types applied to the New Zealand region *Int. J. Climatol.* **14** 711-721
- Kosatsky T 2005 The 2003 European heatwave. *Euro Surveill.* **10**, 148-149
- Lewis J 2018 Kingfish in the harbor climate "sentinels". *Otago Daily Times*. Retrieved from www.odt.co.nz
- Madec G 2008 NEMO the Ocean Engine *Tech. Rep., Notes de l'IPSL*, **27** 1288-1619
- Marshall G J 2003 Trends in the Southern Annular Mode from observations and reanalyses *J. Clim.* **16** 4134-4143
- May P 2004 *Flowering and fruitset in grapevines* Lythrum Press Adelaide
- Menary M B, Kuhlbrodt T, Ridley J, Andrews M B, Dimdore-Miles O B, Deshayes J, Xavier P. et al 2018 Pre-industrial control simulations with HadGEM3-GC3.1 for CMIP6. *Journal of Advances in Modeling Earth Systems*. <https://doi.org/10.1029/2018MS001495>

- 1 Morton J 2018 Marine heatwave changes snapper spawning behaviour. New Zealand Herald. Retrieved
2 from www.nzherald.co.nz
- 3 Mullan A B, Stuart S J, Hadfield M G, Smith MJ 2010 Report on the Review of NIWA's 'Seven-Station'
4 Temperature Series NIWA Information Series No. 78. 175 pp
- 5 Mullan A B, Sood A, and Stuart S 2016 Climate Change Projections for New Zealand: Atmosphere
6 Projections Based on Simulations from the IPCC Fifth Assessment *Wellington: Ministry for the*
7 *Environment*
- 8 Mullan A B, Wratt D S and Renwick J A 2001 Transient model scenarios of climate changes for New
9 Zealand. *Weather and Climate* **21** 3-33
- 10 Olita A 2007 Effects of the 2003 European heatwave on the Central Mediterranean Sea: surface fluxes and
11 the dynamical response. *Ocean Science* **3** (2) 273-289
- 12 Oliver E C J, Benthuyesen J A, Bindoff N L, Hobday A J, Holbrook N J, Mundy C N and Perkins-Kirkpatrick SE 2017 The unprecedented
13 2015/16 Tasman Sea marine heatwave. *Nature Communications* **8** doi:10.1038/ncomms16101
- 14 Oliver E C J, Benthuyesen J A, Bindoff N L, Hobday A J, Holbrook N J, Mundy C N and Perkins-Kirkpatrick.
15 SE 2017 The unprecedented 2015/16 Tasman Sea marine heatwave. *Nature Communications* **8**
16 doi:10.1038/ncomms16101
- 17 Oliver E C J, Donat M G, Moore P J, Smale D A, Alexander L V, Benthuyesen J A, Feng M, Gupta A S,
18 Hobday A J, Holbrook N J, Perkins-Kirkpatrick S E, Straub S C and Wernberg T 2018 Longer and
19 more frequent marine heatwaves over the past century. *Nature Communications* **9**, 1324.
20 doi:10.1038/s41467-018-03732-9
- 21 Otto F E L, Massey N, van Oldenborgh G J, Jones R G, Allen M R 2012 Reconciling two approaches to
22 attribution of the 2010 Russian heat wave *Geophys. Res. Lett.* **39** L04702
- 23 Perkins S E and Alexander L V On the Measurement of Heat Waves *J. Clim.* **26**
24 4500–17 doi.org/10.1175/JCLI-D-12-00383.1
- 25 Parker A, Garcia de Cortazar-Atauri I, Chuine I, Barbeau G, Bois B, Boursiquot J-M, Cahurel J-Y, Clavierie
26 M, Dufourcq T, Geny L, Guimberteau G, Hofmann RW, Jacquet O, Lacombe T, Monamy C, Ojeda
27 H, Panigai L, Payan J-C, Lovelle BR, Rouchaud E, Schneider C, Spring J-L, Storchi P, Tomasi D,
28 Trambouze W, Trought M and van Leeuwen C 2011 Classification of varieties for their timing of
29 flowering and véraison using a modelling approach: A case study for the grapevine species *Vitis*
30 *vinifera* L. *Agricultural and Forest Meteorology* **180** 249-264
- 31 Parker A K 2013 Modelling phenology and maturation of the grapevine *Vitis vinifera* L.: Varietal differences
32 and the role of leaf area to fruit weight ratio manipulations *Lincoln University PhD thesis* Lincoln
33 University
- 34 Perkins S E and Alexander L V 2013 On the measurement of heatwaves *J. Climate* **25** 4500 - 4517
- 35 Perkins-Kirkpatrick, S. E., King, A. D., Couston, E. A., Grosese, M. R., Oliver, E. C. J., Holbrook, N. J.,
36 Lewis, S. C. and Pourasghar, F. 2018 The role of natural variability and anthropogenic climate
37 change in the 2017/18 Tasman Sea marine heatwave. *Bulletin of American Meteorological Society*
38 doi:10.1175/BAMS-D-18-0116.1
- 39 Poloczanska ES, Burrows M T, Brown C J, Molinos J C, Halpern B S, Hoegh-Guldberg O, Kappel C V,
40 Moore P J, Richardson A J, Schoeman D S and Sydeman W J 2016, Responses of Marine
41 Organisms to Climate Change across Oceans, *Front. Mar. Sci* **3** 62 doi:10.3389/fmars.2016.00062.
- 42 Reynolds R W *et al* 2007 Daily high-resolution-blended analyses for sea surface temperature *J. Clim.* **20**
43 5473–596
- 44 Ridgway K R, Dunn J R and Wilkin J L 2002 Ocean interpolation by four-dimensional least squares -
45 Application to the waters around Australia *J. Atmos. Ocean. Tech.* **19** (9) 1357-1375
- 46 Roemmich, D., & the Argo Steering Team. 2009. Argo: The challenge of continuing 10 years of progress.
47 *Oceanography*, 22(3), 46-55. doi:http://dx.doi.org/10.5670/oceanog.2009.65
- 48 Saha S, Nadiga S, Thiaw C, Wang J, Wang W, Zhang Q, Van den Dool H M, Pan H L, Moorthi S, Behringer
49 D, Stokes D, Pena M, Lord. S, White G, Ebisuzki W, Peng P and, Xie P. 2006 The NCEP climate
50 forecast system *J. Climate* **19** 3483–3517
- 51 Salinger M J 1979 New Zealand climate: the temperature record, historical data and some agricultural
52 implications *Climatic Change* **2** 109-126
- 53 Salinger M J 1980 The New Zealand temperature series *Climate Monitor* **9** (4) 112-118
- 54 Salinger M J, Hobday A J, Matear R J, O'Kane T J, Risbey J S, Eveson J P, Fulton, E A, Feng M, Plaganyi
55 E.E, Poloczanska E S, Marshall A G and Thompson P A 2016 Decadal-Scale Forecasting of
56 Climate Drivers for Marine Applications. *Advances in Marine Biology*
57 doi:10.1016/bs.amb.2016.04.002
- 58 Schär C, Vidale P L, Lüthi D, Frei C, Häberli C, Liniger M A and Appenzeller C 2004 The role of increasing
59 temperature variability in European summer heatwaves *Nature* **427** 332-336
- 60 Sirguey P, Still H, Cullen N J, Dumont M, Arnaud Y and Conway J P 2016 Reconstructing the mass balance
of Brewster Glacier, New Zealand, using MODIS-derived glacier-wide albedo *The Cryosphere* **10**
2465-2484 <https://doi.org/10.5194/tc-10-2465-2016>

- 1 Sparnocchia S, Schiano M E, Picco P, Bozzano R and Cappelletti A 2006 The anomalous warming of
2 summer 2003 in the surface layer of the Central Ligurian Sea (Western Mediterranean) *Ann.*
3 *Geophys.* **24** 443–452
- 4 Storkey D, Blaker A T, Mathiot P, Megann A, Aksenov Y, Blockley E W, Calvert D, Graham T, Hewitt H T,
5 Hyder P, Kuhlbrodt T, Rae J G L and Sinha B 2018 UK Global Ocean GO6 and GO7: A traceable
6 hierarchy of model resolutions, *Geosci. Model Dev.* **11(8)** 3187–3213, doi:10.5194/gmd-11-3187-
7 2018.
- 8 Tait A, Henderson R, Turner R and Zheng X G 2006 Thin plate smoothing spline interpolation of daily rainfall
9 for New Zealand using a climatological rainfall surface *Int. J. Climatol.*, **26** 2097-2115
- 10 Trenberth K E 1976 Fluctuations and trends in indices of the southern hemispheric circulation, *Quart. J. Roy.*
11 *Met. Soc.* **102** (431) 65-75
- 12 Trought M, Parker A, Sturman A and Agnew R 2016 Changing New Zealand climate equals a changing New
13 Zealand terroir? In: Jones GV, Doran N eds. 11th International Terroir Congress, McMinnville,
14 Oregon, USA 59-64
- 15 Troup A J 1965 The 'southern oscillation'. *Quart. J. Roy. Met. Soc.* **91** (390) 490-506
16 doi.org/10.1002/qj.49709139009
- 17 Tsujino H, Urakawa , Nakano , Small , Kim , Yeager , Danabasoglu G, Suzuki T, Bamber J L, Bentsen M,
18 Boening C, Bozec A, Chassignet E, Curchitser E, Dias F B, Durack P J, Griffies S M, Harada Y,
19 Ilicak M, Josey S, Kobayashi C, Kobayashi S, Komuro Y, Large W G, Le Sommer J, Marsland S,
20 Masina S, Scheiner M, Tomita H, Valdivieso M and Yamazaki D 2018
21 input4MIPs.CMIP6.OMIP.MRI.MRI-JRA55-do-1-3. Version YYYYMMDD^[1]. Earth System Grid
22 Federation. <http://doi.org/10.22033/ESGF/input4MIPs.2205>.
- 23 van Vuuren D P, Edmonds J, Kainuma M, Riahi K, Thomson A, Hibbard K, Hurtt G C, Kram T, Krey V,
24 Lamarque J F, Masui T, Meinshausen M, Nakicenovic N, Smith S and Rose S K 2011 The
25 representative concentration pathways: an overview. *Climatic Change* **109** 5–31
- 26 Williams J, Morgenstern O, Varma V, Behrens E, Hayek W, Oliver H, Dean S, Mullan A B and Frame D
27 2016 Development of the New Zealand Earth System Model: NZESM *Weather Clim.* **36** 25–44
- 28 Willsman A, Chinn T, Macara G 2018 New Zealand Glacier Monitoring: End of summer snowline survey
29 2017. New Zealand Ministry of Business, Innovation and Employment , Report 2018176EI, 149 pp.
- 30 Wernberg T, Bennett S, Babcock R C, de Bettignies T, Cur, K, Depczynski M *et al* 2016 Climate-driven
31 regime shift of a temperate marine ecosystem *Science* **353** (6295) 169–172
32 <http://doi.org/10.1126/science.aad8745>
- 33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig. 1. The 2017/18 New Zealand atmosphere and ocean heatwaves. **a.** The New Zealand air temperature (NZT) series 1909 – 2018 summer (DJF) anomaly compared with the 1981 – 2010 normal. The red curve is a 15-year filtered average, and black dotted line the trend. **Inset:** Mean NZT August 2017 – April 2018 anomalies. **b.** Summer (DJF) indices of climate extremes for New Zealand from station data and VCSN. Station indices are warm days in red (percentage of time when daily maximum temperature > 90th percentile), warm nights in blue (percentage of time when daily minimum temperature > 90th percentile). The VCSN summer (DJF) count when daily maximum temperature $\geq 25^{\circ}\text{C}$ (in black). **c.** Summer (DJF) SST anomalies 1900-2018 (ERSSTv5) averaged over the Tasman Sea region (26°S – 46°S , 150°E – 174°E) using 1961 – 1990 climatology (Bureau of Meteorology, Australia). **d.** Composite December 2017 – January 2018 SST anomalies (from NOAA ERSSTv5) using the 1981 – 2010 normal with contours at 0.5°C intervals. **e-j.** First row: Time series of summer (Dec-Feb) sea surface temperature (SST) climatology (1981-2011; blue), 90th percentile MHW threshold (orange) and summer 2017/18 SST time series (black) from **(e)** the eastern Tasman Sea (160 - 175°E , 36 - 48°S) and **(f)** the Portobello Marine Laboratory (45.88°S , 170.50°E). The red shaded regions identify periods associated with MHWs detected in the SST time series from each location using the Hobday et al. (2016) definition. Second row: The duration of each MHW detected in the SST time series for **(g)** the eastern Tasman Sea and **(h)** Portobello Marine Laboratory. The red shaded bar highlights MHWs detected during summer 2017/18. Third row: As above but showing the maximum intensity of each MHW detected in the SST time series for **(i)** the eastern Tasman Sea and **(j)** Portobello Marine Laboratory.

Fig. 2. Atmospheric pressure anomalies, tropical cyclone occurrence and sub surface temperature anomalies. **a.** NCEP December 2017 – February 2018 mean sea level pressure anomalies using the 1981-2010 reference period, contour interval 1hPa, negative contours blue, positive red and zero is black; **b.** Named tropical cyclone (TC) tracks for the southwest Pacific for the 2017/18 austral summer to the end of April 2018 based on the latest SPEArTC dataset (Diamond et al 2012); **c.** TC frequency of occurrence anomalies (right panel) calculated as the differences from the 1981–2010 climatology from SPEArTC from those TC seasons, contour interval is 0.5 TC occurrences; red contours positive; blue contours negative; zero contour omitted; **d.** December 2017 – February 2018 sub-surface temperature anomalies for the Pacific Ocean from 40 – 45°S using the 1981-2010 reference period for Global Ocean Data Assimilation System (GODAS) sub-surface temperatures; **e.** Regionally-averaged ARGO float measurements for the eastern Tasman Sea (35 - 45°S , 160 – 172.5°E) of the mean temperature anomaly over the period January 2017 – May 2018.


Fig.3. Ocean modelling: Driving mechanisms for the MHW. Monthly mean values for wind speed anomaly (a-c), MHW intensity (d-f), and mixed layer depths anomaly (g-i) for the period November 2017 – January 2018 based on the model hindcast. To identify MHWs the Hobday et al. (2016) definition has been used. The light blue dashed box identifies the region which has been used to compute area-averages for timeseries in (j) and (k). Annual NDJ (Nov-Dec-Jan) area averaged (Tasman box) anomalies for surface heat fluxes (W/m^2) and wind speeds (m/s) are presented in (j) and SST (C) and thermocline depth (m) in (k). Mean climatological values for NDJ are given in parentheses.

Fig. 4. Southern Alps ice volume and seasonal snow. **a.** Southern Alps ice volume change (km^3 of water equivalent), between years, for all glaciers of the Southern Alps from 1962 to 2018. **b.** Mean (blue) MODIS glacier-wide albedo (2000-2013) and the 2017/18 season (black). Light shading indicates the maximum and minimum range, and dark shading the 25 to 75 percentile range. **c.** estimated water stored as seasonal snow (mm) from SnowSim for the period 1987 – 2016. The year from 1 Apr 2017 – 3 Mar 2018 is shown as the blue line.

Fig. 5. Marlborough Sauvignon Blanc grapevine maturity modelling. **a.** Degree day anomaly from the long-term average accumulated degree days (base 0°C) from 29 August. Estimated flowering (F), véraison (V) and 20°Brix (20) dates using the Grapevine Flowering and Véraison (GFV) model. The date of the phenological events F, V and 20, reflects the degree day anomaly (i.e. the greater the anomaly, the earlier phenology – diagonal lines). The harvest year is the latter year from the austral spring to summer period. **b.** Mean daily temperature from estimated véraison to estimated 20°Brix (maturation (GFV modelled values) for Marlborough, New Zealand for harvest years 1947 - 2018.

Fig.6. Schematic of likely driving mechanisms for the 2017/18 summer New Zealand ocean and atmosphere heatwave.

Figure 1.






Figure 2.




Figure 3.





Figure 4.

**Figure. 5**




Figure 6.

Accepted Manuscript

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60